

# Seasonal dynamics of the larval distribution and settlement of the clam *Ruditapes philippinarum* in the Suo-Nada Sea, Japan

Naoaki Tezuka, Masami Hamaguchi, Manabu Shimizu, Hideki Iwano, Takahiko Tawaratsumida, Shigeru Taga

Coastal Ecosystems, 2016, vol 3, 1-15





© 2016 Society for Coastal Ecosystems Studies + Asia Pacific

Regular Paper



# Seasonal dynamics of the larval distribution and settlement of the clam *Ruditapes philippinarum* in the Suo-Nada Sea, Japan

Naoaki Tezuka<sup>1</sup>, Masami Hamaguchi<sup>1</sup>, Manabu Shimizu<sup>2</sup>, Hideki Iwano<sup>3</sup>, Takahiko Tawaratsumida<sup>4</sup>, Shigeru Taga<sup>5</sup>

<sup>1</sup>National Research Institute of Fisheries and Environment of Inland Sea, 2-17-5 Maruishi, Hatsukaichi, Hiroshima 739-0452, Japan

<sup>2</sup>National Research Institute of Fisheries Science, 2-12-4 Fukuura, Kanazawa, Yokohama, Kanagawa 236-8648
<sup>3</sup>Oita Prefectural Agriculture, Forestry and Fisheries Research Center, Fisheries Research Division, 3386 Kuresaki, Bungo-Takada, Oita 879-0617

<sup>4</sup>Fukuoka Fisheries and Marine Technology Research Center, Buzen Sea Laboratory, 76-30 Unoshima, Buzen, Fukuoka 828-0022

<sup>5</sup>Yamaguchi Prefectural Fisheries Research Center, Naikai Fisheries Research Division, 437-77 Aio-Futajima, Yamaguchi 754-0893

Corresponding author : N. Tezuka, e-mail: tezukan9@fra.affrc.go.jp

#### Abstract

Understanding the causes of spatiotemporal variations in the scale of larval settlement is important for population dynamics studies in bivalves. This study investigated the seasonal abundance of the asari or Manila clam (Ruditapes philippinarum) larvae over a 4-year period (2004–2007) and their settlers over 3 years (2005–2007) in the Suo-Nada Sea, Japan. Seasonal differences in larval transport were examined by numerical simulation using 3D ocean-modelling. During the 2004-2007 spawning seasons, larval numbers peaked two or three times in June/July, August/September, and October/November. Settler occurrence was uncoupled with larval occurrence; settler density was >10 times higher in October/November than in other months. Numerical simulation suggested that the extent of larval transport differed seasonally; larval transport via loss from the Suo-Nada Sea was estimated to be 20% in June/July, whereas it was almost 0% in November. However, this could not explain the seasonal difference (>10-fold) in settler density. In addition, the density of larger larvae (>180 µm) averaged over the years 2004–2007 was higher in June/July than in October/ November, as a result of spawning and larval loss (via transport and mortality), suggesting that larval supply alone could not explain the seasonal differences in settler density. These results suggest that the seasonal differences in settler density were affected more by variation in mortality during the settlement and/or early post-settlement stages, which may depend on environmental conditions at the settling site, rather than by larval supply.

Keywords: clam, larval supply, larval transport, settlement

#### Introduction

For many marine bivalves including the asari or Manila clam *Ruditapes philippinarum* (Adams & Reeve 1850), spatio– temporal differences in larval supply constitute a potential cause of variations in the scale of larval settlement (Young *et al.* 1998; Shanks & Brink 2005). Understanding the causes of variations in larval supply and settlement, along with post-settlement processes, is important for studies on the population dynamics of bivalves (Hunt & Scheibling 1997; Pineda, Hare & Sponaugle 2007).

*R. philippinarum* was originally distributed in the temperate to subarctic regions along the east coast of the Pacific Ocean, but is currently found in many other areas worldwide because of its introduction for aquaculture (Goulletquer 1997). Population studies on *R. philippinarum* have attracted attention because of its commercial and ecological importance, especially after a population decline in Japan (Miyawaki & Sekiguchi 1999; Ishii *et al.* 2001; Toba

*et al.*2007; Tamaki *et al.* 2008; Tezuka *et al.* 2012). Spawning and larval occurrence of the asari clam has been observed during spring to autumn at water temperatures above 14 °C (Miyawaki & Sekiguchi 1999; Ishii *et al.* 2001; Matsumura *et al.* 2001; Drummond, Mulcahy & Culloty 2006). Dense larval settlement has often been recorded in autumn in temperate regions of Japan (Miyawaki & Sekiguchi 1999; Ishii *et al.* 2001; Toba *et al.* 2007). However, the causes of seasonal variation in settler density and the decoupling between larval occurrence and settlement are as yet unclear.

The Suo-Nada Sea was renowned as one of the major *R. philippinarum* fishery ground in Japan until the population drastically declined in the late 1980s (see Tezuka *et al.* 2012, 2014). Surveys conducted in the 1970s and early 1980s, before the population decline, showed that larval abundance peaked in spring and autumn, and that settlers were observed mainly in autumn (Inoue 1980; Fujimoto *et al.* 1985). No larval surveys have been carried out after the population decline; however, seasonal observations of



**Fig. 1** (a)-(b), Map of the Suo-Nada Sea, Japan. Dashed line in (b) is the boundary used to calculate the larval retention rate (see text). (c)-(d), Nakatsu tidal flats where larval settlement was recorded.

- 2 -

settlement on the Nakatsu tidal flats in the Suo-Nada Sea have been reported recently by Tezuka *et al.* (2012), and settlers have been observed in autumn.

To ascertain the causes of seasonal differences in settler density observed on the Nakatsu tidal flats during 2005–2007 by Tezuka *et al.* (2012), we investigated the abundance and distribution of *R. philippinarum* larvae between April and November over the period 2004–2007. In addition, seasonal

differences in larval transport were examined by numerical simulation. Data for settler numbers were obtained from the Tezuka *et al.* (2012) study and reanalyzed to determine the periods of larval settlement using shell-length distribution. The study will increase our understanding of the causes of seasonal differences in settler density and larval numbers after the clam population decline in the Suo-Nada Sea.



**Fig. 2** Modelled larval particles were set in 9 areas along the Suo-Nada Sea coast for numerical simulation of *Ruditapes philippinarum* larval transport. A total of 1000 modeled larval particles were set on 10-model grids in each area. The model grids in each area were indicated as "+" symbols with identical colour.

#### Materials and Methods

#### Sampling of larvae and settlers

Sampling of *R. philippinarum* larvae was conducted in the Suo-Nada Sea between April and November over the period 2004–2007 (Fig. 1). Larvae were collected by pumping 200 L of seawater from 5 m depth and screening onto a 50-µm mesh net. The sampling depth was changed to an intermediate level in cases where the sampling stations were shallower than 10 m. In the periods 19-22 June 2007, 20-23 August 2007, 15-19 October 2007, and 16-19 November 2007, multilayer sampling from 2, 5 and 10 m depths was conducted to ascertain the vertical distribution of clam larvae. Samples were frozen until needed for identification, enumeration and shell length measurement. After thawing, larvae were identified using fluorescent antibodies and counted under a fluorescence microscope (Matsumura et al. 2001; Kasuya, Hamaguchi & Furukawa 2004). Shell length was measured using an evepiece to the nearest 10 µm, and up to 100 individuals were measured per sample.

*R. philippinarum* settlers were collected at two stations on the Nakatsu tidal flats during the period 2005–2007 (Fig. 1; see Tezuka *et al.* 2012) by taking three, 1-cm deep surface-

sediment samples with a core sampler (40 mm in diameter). Samples were preserved by freezing until settlers were counted and shell lengths measured to the nearest 0.1 mm.

Numerical simulation of larval transport

A 3D numerical ocean model was used for simulation of *R. philippinarum* larval transport in the Suo-Nada Sea. Particletracking experiments were conducted using the same flow-field reported by Chang *et al.* (2009). A total of 1000 modelled larval particles were set in each of nine areas along the Suo-Nada coast (Fig. 2) and tracked by the Euler– Lagrange method for 14 days (1000 particles per 10-model grids per area). The simulation was conducted over two seasonal periods: June/July (from June 30 to July 17) and November (from November 7 to November 27).

The position of the particle  $X_{n+1}(x^{n+1}, y^{n+1})$  at time n+1, which was  $X_n(x^n, y^n)$  at time n, can be calculated by the following equation:

$$X_{n+1} = X_n + V\Delta t + (\partial V/\partial t + V\nabla V)\Delta t^2/2 + R$$

where *V* denotes the horizontal velocity vector of residual flow;  $\Delta t$ , the time step;  $\nabla$ , the horizontal gradient. *R* is the

\_\_\_\_\_

dispersion due to turbulence and is given by the following equation:

$$R = \gamma (2 \Delta t D_h)^{1/2}$$

where  $\gamma$  is the normal random number whose average is zero and whose standard deviation is 1.0.  $D_h$  is the horizontal dispersion coefficient that depends on Smagorinsky diffusivity (Mellor 2004), which varies with horizontal shear stress.

Vertical migration of larval particles was hypothesized as larvae were located at 3 m depth from day 0 to day 11 but, then moved to 1 m above the ocean floor between day 12 and day 14. This assumption for the vertical migration of larval particles is based on observational studies on the vertical distribution of larvae, including this study (see Results, Fig. 6), which showed that smaller larvae (D- and umbo-shaped) were found at ~3 m, whereas larger larvae (settling larvae) found in the bottom layer (Suzuki *et al.* 2002; Ishii, Sekiguchi & Jinnai 2005; Kuroda 2005; Toba *et al.* 2012; Bidegain *et al.* 2013). The larval stage was assumed to last 2 weeks (14 days) in this study, although it can vary with temperature and food availability (Helm & Bourne 2004).

The larval retention rates, i.e. the ratio of larval particles remaining within the Suo-Nada Sea, after the 14-day

simulation were calculated as follows:

Retention rate after 14 days =  $100 \times P_a / P_o$ 

where  $P_a$  is the number of particles remaining within the Suo-Nada Sea after the 14-day simulation, and  $P_o$  is the number of particles within the Suo-Nada Sea on Day 0. The boundary of the Suo-Nada Sea was set on a line through two points (131.7°E, 33.7°N) and (132.0°E, 34.0°N) (see Fig. 1), and particles located to the western side of the boundary line were treated as inside the Suo-Nada Sea. Retention rates were calculated for each of the nine areas where particles were released for two seasonal periods.

Seasonal difference in larval density by size

The density of settling larvae, i.e., the larval supply, could vary with spawning biomass and larval loss due to transport and mortality. If we assume that larval loss is very high, including loss via transport and death, larger larvae are unlikely be observed unless spawning is very intense. To examine the seasonal differences in the density of settling larvae, the larval densities were compared between June/July and October/November, using the combined data for 2004–2007.



**Fig. 3** (a) Seasonal dynamics of *Ruditapes philippinarum* larval distribution in the Suo-Nada Sea in 2004.







Fig. 3 (c) Seasonal dynamics of *Ruditapes philippinarum* larval distribution in the Suo-Nada Sea in 2006.



**Fig. 3** (d) Seasonal dynamics of *Ruditapes philippinarum* larval distribution in the Suo-Nada Sea in 2007.

#### Results

#### Seasonal dynamics of larval distribution and settlement

Distribution of *R. philippinarum* larvae from April to November 2004–2007 is shown in Fig. 3a-d. Seasonal changes in planktonic larval abundance are shown in Fig. 4. *R. philippinarum* larvae were observed sporadically from April to November, with peak numbers being recorded two or three times in a year, i.e. June/July, August/September and October/November (Figs. 3 and 4). Larvae were distributed more widely in the Suo-Nada Sea during June/July (e.g. 13–16 June 2005, 19–2 June 2006 and 19–22 June 2007) than in October/November (e.g. 20–23 November 2005, 30 October–2 November 2006 and 15–19 October 2007) (Fig. 3). The maximum recorded planktonic larval density was c. 2000 individuals m<sup>-3</sup>, observed in November 2006 and in June 2007 (Fig. 4).

The size-frequency distribution of *R. philippinarum* settlers on the Nakatsu tidal flats between May–December 2005–2007 is shown in Fig. 5. New settlers (< 0.5 mm in shell length) were found almost exclusively in October/



Fig. 4 Larval density of *Ruditapes philippinarum* in the Suo-Nada Sea during April–November of 2004–2007.



**Fig. 5** Size-frequency distributions of *Ruditapes philippinarum* settlers on the Nakatsu tidal flats, in the Suo-Nada Sea, during May–December 2005–2007. Data from two stations were combined.

 Table 1. Retention rate (%) of particles within the Suo-Nada Sea after a 14-day simulation. Retention rates were calculated for each of the 9 areas where particles were released over two seasonal periods, June/July and November.

	Area								moon	
	1	2	3	4	5	6	7	8	9	mean
June/July	85.4	93.7	40.4	32.2	100	100	75.6	89.9	98.7	79.5
Nov	99.7	100	100	100	100	100	100	100	99.4	99.9

November for the 3-year period (Fig. 5). The density of new settlers (< 0.5 mm in shell length) differed by more than 10-fold between summer and autumn: 200-1000 individuals  $m^{-2}$  in June–August, and 4000-13,000 individuals  $m^{-2}$  in October/November.

#### Vertical distribution of larvae

Vertical distribution of *R. philippinarum* larvae, inferred from the multilayer sampling trials, are shown in Fig. 6. During 19–22 June 2007, smaller larvae (120–180  $\mu$ m in shell length) were more abundant at the 2 and 5 m depths than at 10 m, whereas larger larvae (>180  $\mu$ m) were more abundant at 10 m. During the other time-periods, larvae were less abundant than during 19–22 June 2007, being found almost exclusively at 2 m.

#### Numerical simulation of larval transport

Results of the numerical simulation of larval particle-tracking are shown in Fig. 7. In the June/July period, released larval particles were carried extensively by two major currents that developed in the Suo-Nada Sea; a circular current in the central region and a northward current along the western coast. In the November simulation, the currents had weakened and the larval particles were retained within the Suo-Nada Sea. The retention rates for particles within the Suo-Nada Sea after the 14-day simulations for June/July and November are shown in Table 1. In the June/July period, retention rates of larval particles varied by area where the particles were released, with rates ranging from 32.2% (Area 4) to 100% (Areas 5 and 6). On average, 79.5% of the particles were retained within the Suo-nada Sea, while the



**Fig. 6** Abundance of *Ruditapes philippinarum* larvae at three different depths in the Suo-Nada Sea in (a) 19–22 June, (b) 20–23 August, (c) 15-19 October, and (d) 16–19 November, all in 2007.

7day [7 Jul.]		131°00' 20' 40' 132°00'	7day [14 Nov.]	131°00 200 400 132°00		ylune/July
5day [5 Jul.]		7 131°00 20 40 132°00 14day [14 Jul.]	5day [12 Nov.]	7 131°0″ 20 40 132°0″		or 131°00' 20' 40' 132°00' art in the Suo-Nada Sea, for (a) <b>.</b>
3day [3 Jul.]		7 131°00 20 40 132°0 13day [13 Juli]	3day [10 Nov.]	y 131°0° 20° 40° 132°0		or 131°00' 20' 40' 132°0 per philippinarum larval transpc deled larval particles released f
[.lu[ 1] 1day		131°00 20° 40° 132°00 12day [12 Jul.]	1day [8 Nov.]			r 131'00' 20' 40' 132'0 numerical simulation of <i>Rudita</i>
(a) June-July Initial [30 Jun.]	34°00	131:00 20 40 132:00 9day [9 Jul.] 33:40 33:40 131:00 20 40 132:00	(b) November Initial [7 Nov.]		34°00 I I NOV.J	33°40 <sup>(1)</sup> 131°00 <sup>(2)</sup> 20 <sup>(4)</sup> 132°0 <b>Fig. 7</b> Results of the and (b) November. Idei

rest were lost by being transported from the Suo-Nada Sea in the June/July period. For the November period, retention rates were higher than in the June/July period at >99%.

#### Seasonal difference in larval density by size

The densities of planktonic larvae of different sizes are compared between June/July and October/November for combined data (2004–2007) (Fig. 8). The mean larval density of all combined size classes was 79.2 individuals m<sup>-3</sup> in June/July and 36.3 individuals m<sup>-3</sup> in October/November, i.e., approximately twice as higher in the former. Of these, larger larvae (>180 µm) constituted 8.0% in June/July (6.3 individuals m<sup>-3</sup>) and 10.2% in October/November (3.7 individuals m<sup>-3</sup>).



2004–2007. Bars indicate standard errors.

**Fig. 8** Mean density of larvae in different size classes in June/July (upper) and October/November (lower),

#### Discussion

This study reported the seasonal dynamics of *R. philippinarum* larval distribution in the Suo-Nada Sea and the number of settlers on the Nakatsu tidal flats. Larval occurrence and settlement events were not necessarily coupled and, although larvae were recorded from April to November, settlement occurred almost exclusively in October/November. Numerical simulation suggested that larval loss via current transport was higher in June/July than in November. However, the actual density of larger larvae (>180  $\mu$ m) was higher in June/July than in October/November that the supply of larvae was not the cause of seasonal differences in settler density.

The seasonal occurrence pattern of larvae and settlers observed in this study was similar to that reported in previous studies within the Suo-Nada Sea and other regions. Inoue (1980) reported that planktonic larval density peaked in May and October/November in 1976 on the northern coast of the Suo-Nada Sea (maximum larval density was 14,000 individuals m<sup>-3</sup> in October 1976), and that new settlers were found in abundance (26,700 individuals m<sup>-2</sup> in November 1976) in October/November, but very few, new settlers were

recorded in other months. Fujimoto et al. (1985) observed that larval density peaked in April, June/July, September and November during the period 1983-1984 on the southwest coast of the Suo-Nada Sea, with a maximum density of 4550 individuals m<sup>-3</sup> in November 1983 and 6140 individuals m<sup>-3</sup> in April 1984 (larval density was calculated as the mean from eight sampling stations). An increase in new settlers was observed in October/November 1983 (maximum of 350,000 individuals m<sup>-2</sup> in November 1983) and in November/ December 1984 (max, 20,000 individuals m<sup>-2</sup>). Although the maximum larval density observed in the current study was lower than those recorded in the 1970s and 1980s, the seasonal pattern was consistent among these studies. The difference in maximum larval density could be a result of differences in the spawning biomass between the 2000s and the 1970–1980s; clam production in the Suo-Nada Sea was 14,800 metric tonnes in 1976, 19,500-31,700 metric tonnes in 1983/1984, and 94-780 metric tonnes during 2005-2007 (Ministry of Agriculture, Forestry and Fisheries, Japan, 2012). Dense larval settlement in autumn has often been observed in other temperate regions of Japan (Miyawaki & Sekiguchi 1999; Ishii et al. 2001; Toba et al. 2007).

The results of numerical simulation suggest that larval

retention in the Suo-Nada Sea should be higher, i.e., lower larval loss rates, in October/November than June/July, owing to seasonal differences in the strength of two major currents in the Suo-Nada Sea. As stratification developed in June/ July, a circular current developed in the central region and a northern current developed along the southwest coast of the Suo-Nada Sea. However, as stratification declined in October/November, the currents weakened. The simulation suggested that larval loss via transport from the Suo-Nada Sea was 20% in June/July and 0% in October/November. However, this disparity could not explain the >10-fold difference in settler density between October/November and June/August. Moreover, averaged over the years 2004–2007, the mean density of >180 µm larvae was higher in June/July than in October/November. This suggests that, even if the larval loss via transport may be higher in June/ July, larval supply, as a result of spawning and larval loss via transport and mortality, was not a limiting factor in June/July under the prevailing conditions.

Factors influencing settler density, other than larval supply, need to be considered, and although not all factors were investigated in this study, several hypotheses may be proposed. First, mortality during the critical period from settling to early post-settlement affects settler density (Hunt & Scheibling 1997), and this mortality may be lower in October/November than in other months. Predation and/ or other factors, such as high temperature in summer, may cause higher mortality in settling larvae or early settlers during months other than October/November. Second, food availability for larvae may differ seasonally, resulting in differences in mortality during settlement (Laing 1995; Tezuka et al. 2009). Third, the rate of larval settlement may vary due to seasonal differences in physical conditions at settling sites, e.g., variation in sediment grain size (Tezuka et al. 2013). If a suitable substrate is not found, larvae could delay settlement (Coon, Fitt & Bonar 1990) and, as a result, become more prone to predation by benthic filterfeeders while drifting near the bottom (Pineda et al. 2010). This study concentrated solely on larval settlement on the tidal flats and other possible sites, e.g., subtidal areas, were not investigated. As such, settlement failure on the tidal flats in seasons other than October/November needs to be investigated further.

#### Acknowledgments

This research was funded by the Fisheries Agency, Japan. We thank Professor Takeoka and Professor Guo from the Center for Marine Environmental Studies at Ehime University for providing the ocean model outputs. We also thank reviewers and the editor for their constructive and valuable comments, which helped us to improve the manuscript.

#### References

- Bidegain G, Bárcena JF, García A & Juanes JA (2013) LARVAHS: Predicting clam larval dispersal and recruitment using habitat suitability-based particle tracking model. Ecological Modelling 268, 78–92.
- Chang P-H, Guo X & Takeoka H (2009) A Numerical Study of the Seasonal Circulation in the Seto Inland Sea, Japan. Journal of Oceanography 65, 721–736.
- Coon SL, Fitt WK & Bonar DB (1990) Competence and delay metamorphosis in the Pacific oyster *Crassostrea gigas*. Marine Biology 106, 379–387.
- Dethier MN, Ruesink J, Berry H & Sprenger AG (2012) Decoupling of recruitment from adult clam assemblages along an estuarine shoreline. Journal of Experimental Marine Biology and Ecology 422-423, 48–54.
- Drummond L, Mulcahy M & Culloty S (2006). The reproductive biology of the Manila clam, *Ruditapes philippinarum*, from the North-West of Ireland. Aquaculture 254, 326–340.
- Fujimoto T, Nakamura M, Kobayashi M, Hayashi I, Takiguchi K, Oda K & Ushima H (1985) On the formation of the asari clam fishery ground. Annual Report of Fukuoka Prefecture Buzen Fisheries Experimental Station S58, pp. 34–106. (In Japanese.)
- Goulletquer P (1997) A Bibliography of the Manila Clam *Tapes philippinarum*. IFREMER, La Tremblade, France. (RIDRV-97.021RN, 122 pp.)
- Helm MM & Bourne N (2004) Hatchery culture of bivalves. A practical manual. Lovatelli A (ed), FAO Fisheries Technical Paper 471. Rome, FAO. 177pp.

- Hunt HL & Scheibling RE (1997) Role of early post-settlement mortality in recruitment of benthic marine invertebrates. Marine Ecology Progress Series 155, 269–301.
- Inoue Y (1980) On the ecology of the asari clam and the environment in Omi- and Yamaguchi-wan sea. Fisheries Engineering 16, 29–35. (In Japanese.)
- Ishii R, Sekiguchi H, Nakahara Y & Jinnai Y (2001) Larval recruitment of the manila clam *Ruditapes philippinarum* in Ariake Sound, southern Japan. Fisheries Science 67, 579–591.
- Ishii R, Sekiguchi H, Jinnai Y (2005) Vertical distributions of larvae of the clam *Ruditapes philippinarum* and the striped horse mussel *Musculista senhousia* in eastern Ariake Bay, southern Japan. Journal of Oceanography 61, 973–978.
- Kasuya T, Hamaguchi M & Furukawa K (2004) Detailed observation of spatioal abundance of clam larvae *Ruditapes philippinarum* in Tokyo Bay, central Japan. Journal of Oceanography 60, 631–636.
- Kuroda N (2005) Larval transportation and settlement mechanism to tidal flat in Japanese littleneck clam *Ruditapes philippinarum*. Bulletin of Fisheries Research Agency, Supplement 3, 67–77. (In Japanese with English abstract.)
- Laing I (1995) Effect of food supply on oyster spatfall. Aquaculture 131, 315–324.
- Matsumura T, Okamoto S, Kuroda N & Hamaguchi M (2001) Temporal and spatial distributions of planktonic larvae of the clam *Ruditapes philippinarum* in Mikawa Bay; Application of an immunofluorescence identification method. Japanese Journal of Benthology 56, 1–8. (In Japanese with English abstract.)
- Mellor G (2004) Users guide for a three-dimensional, primitive equation, numerical ocean model (the June 2004 revision). Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, NJ 08544-0710, 1–56.
- Ministry of Agriculture, Forestry and Fisheries, Japan (2012) Annual report of statistics on fisheries and aquaculture production in 2010. Nourin Toukei Kyoukai, Tokyo. (In Japanese.)
- Miyawaki D & Sekiguchi H (1999) Interannual variation of bivalve populations on temperate tidal flats. Fisheries Science 65, 817–829.
- Pineda J, Hare JA & Sponaugle S (2007) Larval transport and dispersal in the coastal ocean and consequences for population connectivity. Oceanography 20, 22–39.
- Pineda J, Porri F, Starczak V & Blythe J (2010) Causes of decoupling between larval supply and settlement and consequences for understanding recruitment and population connectivity. Journal of Experimental Marine Biology and Ecology 392, 9–21.

- Shanks AL & Brink L (2005) Upwelling, downwelling, and crossshelf transport of bivalve larvae: test of a hypothesis. Marine Ecology Progress Series 302, 1–12.
- Suzuki T, Ichikawa T & Momoi M (2002) The approach to predict sources of pelagic bivalve larvae supplied to tidal flat areas by receptor mode model: a modeling study conducted in Mikawa Bay. Bulletin of the Japanese Society of Fisheries Oceanography 2, 88–101. (In Japanese with English abstract.)
- Tamaki A, Nakaoka A, Maekawa H & Yamada F (2008) Spatial partitioning between species of the phytoplanktonfeeding guild on an estuarine intertidal sand flat and its implication on habitat carrying capacity. Estuarine, Coastal and Shelf Science 78, 727–738.
- Tezuka N, Ichisaki E, Kanematsu M, Usuki H, Hamaguchi M & Iseki K (2009) Particle retention efficiency of asari clam *Ruditapes philippinarum* Iarvae. Aquatic Biology 6, 281–287.
- Tezuka N, Kamimura S, Hamaguchi M, Saito H, Iwano H, Egashira J, Fukuda Y, Tawaratsumida T, Nagamoto A & Nakagawa K (2012) Settlement, mortality and growth of the asari clam (*Ruditapes philippinarum*) for a collapsed population on a tidal flat in Nakatsu, Japan. Journal of Sea Research 69, 23–35.
- Tezuka N, Kanematsu M, Asami K, Sakiyama K, Hamaguchi M & Usuki H (2013) Effect of salinity and substrate grain size on larval settlement of the asari clam (Manila clam, *Ruditapes philippinarum*). Journal of Experimental Marine Biology and Ecology 439, 108–112.
- Tezuka N, Kanematsu M, Asami K, Nakagawa N, Shigeta T, Uchida M & Usuki H (2014) *Ruditapes philippinarum* mortality and growth under netting treatments in a population-collapsed habitat. Coastal Ecosystems 1, 1–13.
- Toba M, Yamakawa H, Kobayashi Y, Sugiura Y, Honma K & Yamada H (2007) Observations on the maintenance mechanisms of metapopulations, with special reference to the early reproductive process of the Manila clam *Ruditapes philippinarum* (Adams & Reeve) in Tokyo Bay. Journal of Shellfish Research 26, 121–130.
- Toba M, Yamakawa H, Shoji N & Kobayashi Y (2012) Vertical distribution of Manila clam *Ruditapes philippinarum* larvae characterized through year-round observations in Tokyo Bay, Japan. Nippon Suisan Gakkaishi 78, 1135– 1148. (In Japanese with English abstract.)
- Young EF, Bigg GR, Grant A, Walker P & Brown J (1998) A modelling study of environmental influence on bivalve settlement in the Wash, England. Marine Ecology Progress Series 172, 197–214.



Published: 10 February 2016